State: Assam Agriculture Contingency Plan for District: Darrang

1.0 I	District Agriculture profile						
1.1	Agro- Climatic / Ecological Zone						
	Agro Ecological Sub Region (ICAR)	Assam And Bengal Plain, Hot	Assam And Bengal Plain, Hot Subhumid To Humid (Inclusion Of Perhumid) Eco-Region (15.2)				
	Agro- Climatic Region (Planning Commission)	Eastern Himalayan Region (II)					
	Agro Climatic Zone (NARP)*	North Bank Plain Zone (AS-1)	North Bank Plain Zone (AS-1)				
	List all the districts falling under the NARP Zone	Darrang, Udulguri	Darrang, Udulguri				
	Geographic Coordinates of district	Latitude	Longitude	Altitude			
		20°9′N to 26°95′N	91°45′ E to 92°22′ E				
Name and Address of the concerned ZRS/ZARS/RARS/RRTTS KVK, Darrang, Mangaldai(P.O) Darrang (dt), Assam.784125.							
	Mention the KVK located in the district	KVK, Darrang, Mangaldai(P.O) Darrang (dt), Assam.784125					

1.2	Rainfall	Average (mm)	Number of rainy	Normal Onset	Normal cessation
			days		
	SW monsoon (June –Sept)	1254		1 st week of June	4 th week of September
	NE Monsoon (Oct- Dec)	120		1 st week of October	4 th week of November
	Winter (Jan-March)	199		2 nd week of March	-
	Summer (Apr- May)	377		1 st week of April	-
	Annual	1951		-	

1.	Land use pattern	Geographical	Cultivated	Forest	Land under	Permanent	Cultivable	Land under	Barren and	Current	Oth
3	of the district	area	area	area	nonagricultural	pastures	wasteland	misc tree	uncultivable	fallows	er
	(latest statistics)				use			crops and	land		fall
								groves			ows
	Area ('000ha)	320.68	87.277	207.5	12.434	3.105	3.250	4.760	-	2.362	-

1.4	Major Soils	Area ('000ha)	Percent (%) of total
	Sandy loam soils	67.5	47.46
	Clay loam soils	49.9	35.12
	Silty clay loam soils	22.5	15.63

1.5	Agricultural land use	Area ('000ha)	Cropping intensity%
	Net sown area	73.619	180.30%

A	Area sown more than once	59.116
G	Gross cropped area	132.735

1.6 Irrigation	Area ('000 ha)	Percent (%)	
Net irrigated area	26.070 ha	100.00	
Gross irrigated area	-		-
Rainfed area	-		-
Sources if Irrigation	Number	Area ('000 ha)	% area
Canals	-	-	-
Tanks	-	-	-
Open wells	-	-	-
Bore wells	-	-	-
Lift irrigation	3000	7.527	19.97
Other sources	5930	14.967	39.72
Pump sets	6000	15.144	40.31
Micro-irrigation	-	-	-
Groundwater availability and use	No. of blocks	% area	Quality of water
Over exploited			
Critical			
Semi-critical			
Safe			
Wastewater availability and use			

^{*}Over-exploited: groundwater utilization>100%; critical: 90-100%; Semi-critical: 70-90%; Safe:<70% Source: District Agriculture Office, Darrang, Assam

1.7 Area under major field crops & horticulture etc.

1.7	Major field		Area ('000 ha)									
	crops cultivated		Kharif			Rabi Summer						
		Irrigated	Rainfed	Total	Irrigated	Rainfed	Total	Irrigated	Rainfed	Total	Grand total	
	Sali rice		27.729	27.729							27.729	
	Summer rice								23.513	23.513	23.513	

	Horticulture crops - Fruits	Total area (ha)	Irrigated	Rainfed
1.7	Banana	2, 550	-	2,550
	Papaya	385	-	385
	Orange	495	-	495
	Pineapple	526	-	526
	Guava	215	-	215
	Litchi	185	-	185
	Jack fruit	1, 318	-	1,318
	Mango	150	-	150
	Other fruits	550	-	550
	Horticulture crops - Vegetables	Total area	Irrigated	Rainfed
	Total tuber crops (potato, sweet potato, tapioca)	10, 719	-	10, 719
	Total spices (chilli, turmeric, ginger, coriendar,garlic, onion,black pepper)	8, 211	-	8, 211
	Kharif vegetable	16, 975	-	16, 975
	Rabi vegetables	25, 186	-	25, 186
	Medicinal and Aromatic crops	Total area	Irrigated	Rainfed
	Plantation crops	Total area	Irrigated	Rainfed
	Tea	41, 667	41, 667	-
	Areca nut	4, 750	-	4, 750
	Coconut	981	-	981
	Fodder crops	Total area	Irrigated	Rainfed
	Total Fodder crop area			
	Grazing land			

1.8	Livestock	Number ('000)
	Cattle	438.208
	Buffaloes total	122.19

	Commercial dairy farms	0.042		
	Goat	156.893		
	Sheep	39. 006		
	Pig	54.618		
1.9	Poultry			
	Commercial			
	Backyard	1670.912		
1.10	Inland Fisheries	Area (ha)	Yield (t/ha)	Production (tones)
	Brackish water			
	Fresh water			

1.11	Production and Productivity of Major Crops	Kharif		Kharif Rabi Summer			Total		
Field	crops	Production ('000 t)	Productivity (kg/ha)						
	Rice	326	2400	256	3770	98	2400	680	2857
	Toria	-	-	19.10	6.59	-	-	19.10	6.59
	Jute	-	-	-	-	20.0	2060	20.0	2060
	Blackgram	-	-	-	-	2.10	544	2.10	544
	Wheat	-	-	6.73	982	-	-	6.73	982
	Sugarcane	-	-	-	-	307.75	35793	307.75	35793
	Major Horticultural								
	crops								
	Banana	63750	15000	-	-	-	-	63750	15000
	Papaya	64000	32000	-	-	-	-	64000	32000
	Assam lemon	29865	16500	-	-	-	-	29865	16500
	Pineapple	26220	13800	-	-	-	-	26220	13800
	Coconut	30901	6300	-	-	-	-	30901	6300
	Arecanut	92035	15800	-	-	-	-	92035	15800
	Kharif vegetables	114252	115					114252	115
	Rabi vegetables	-		294550	215	-	-	294550	215


1.	.12 Sowing window for 5 major crops (start and end of sowing period)	Rice	Jute	Blackgram	Toria	Wheat
	Kharif-Rainfed	June to July	March to May	Mid August to Mid September		-
	Kharif-Irrigated	-	-	-	-	-
	Rabi-Rainfed	-	-	-	Mid October to	1 st week of November to

				Mid November	2 nd week of December
Rabi-Irrigated	November to	-	-	-	-
	December				


1.13	What is the Major contingency the District is prone to?	Regular	Occasional	None
	Drought		√ (March & October)	
	Flood		√ (August-Sept)	
	Cyclone		√ (March)	
	Hail storm		√ (March-April)	
	Heat wave		√ (June-July)	
	Cold wave		√ (Dec-January)	
	Frost	·		$\sqrt{}$

14	Include Digital maps of the district for	Location map of district with in State Annexure I	Enclosed: Yes
		Mean annual rainfall as Annexure II	Enclosed: Yes
		Soil map as Annexure III	

Annexure-1



Annexure-II Mean Annual Rainfall



2.0 Strategies for weather related contingencies

2.1 Drought

2.1.1 Rainfed situation

Condition				Suggested Contingency measure	s
Early season drought (delayed onset)	Major Farming situation	Crop/ cropping system	Change in Crop/ cropping system	Agronomic measures	Remarks on Implementation
Delay by 2 weeks 3 rd week of June	Medium land	Sali rice – rabi oilseeds/ pulses – boro rice Sali rice – rabi oilseeds/ pulses Sali rice – Wheat Sali rice – rabi vegetables	No change No change No change No change	Decrease the spacing in paddy	Supply of rice seeds through NFSM and other such programme Supply of weeder under RKVY.
Delay by 4 week 1 st week of July	Medium land	Sali rice – rabi oilseeds/ pulses – boro rice Sali rice – rabi oilseeds/ pulses Sali rice – rabi vegetables	Rice: Satyaranjan, Basundhar Rice: Satyaranjan, Basundhar Rice: Satyaranjan, Basundhar	For the Sali varieties, sowing can be done up to mid or last part of July	
Delay by 6 weeks 3 rd week of July	Upland	Sesame/ kharif blackgram/ Greengram/ Pigeonpea	Sesame/ Pigeonpea	Normal sowing of Pigeonpea & sesame can be done Application of sufficient amount of organic manures before sowing Use higher seed rate Thinning to maintain optimam plant population, Mulching with waste materials	Supply of rice seeds through NFSM and other such programme Supply of weeder under RKVY.
	Medium land	Sali rice - boro rice	Sali rice variety like Jaya, IR- 36 and also medium duration variety Satyaranjan and Basundhara Late Sali variety like Monahar Sali, Andrew Sali, Salpona, Prasadbhog etc can be chosen		Supply of rice seeds through NFSM and other such programme Supply of weeder under RKVY.

			Dono mino lileo Emi O Emi 20		
			Boro rice like Eri-9, Eri-28,		
			Eri-30, Joymoti, Kanaklata		
			can be chosen		
Delay by 8 weeks 1 st	Upland	Sali rice – rabi oilseeds/ pulses	Sali rice variety like Jaya, IR-		Supply of rice seeds
week of August			36 and also medium		through NFSM and
			duration variety		other such programme
			Satyaranjan and		Supply of weeder
			Basundhara		under RKVY.
		Sali rice – rabi oilseeds/ pulses	Sali rice variety like Jaya, IR-		
			36 and also medium		
			duration variety		
			Satyaranjan and		
			Basundhara		
		Sali rice – rabi vegetables	Sali rice variety like Jaya, IR-		
			36 and also medium		
			duration variety		
			Satyaranjan and		
			Basundhara		
		SRI - rabi vegetables	Late Sali variety like		
		STEE TWO TYPES CONTROLLED	Monahar Sali, Andrew Sali,		
			Salpona, Prasadbhog etc		
			can be chosen		
	Medium land	Sali rice/ Sesame/kharif blackgram/	i.Use of late Sali varieties-		Supply of rice seeds
	Medium fand		Monuharsali, Andrewsali,		
		Greengram/ pigeonpea			through NFSM and
			Solpona, Prasadbhog,		other such programme
			Gobinbhog		Supply of weeder
			ii.Varieties suitable for both		under RKVY.
			normal & delayed sowing &		
			planting conditions- Luit,		
			Kapilee, Dishang, Dikhow		
			iii.Staggered planting Sali		
			varieties- Prafulla & Gitesh		
	1			1	

Condition Suggested Contingency measures					
Early Season drought	Major Farming	Crop/ Cropping System	Crop management	Soil nutrient & moisture	Remarks on
(Normal onset)	Situation			conservation measures	Implementation
Normal onset followed by	Up land	Sali rice		Application of urea along	Supply of rice
15 to 20 days dry spell				with manure/FYM in the	seeds through
after sowing leading to				nursery, Sprinkling of	NFSM and other

poor germination / crop stand etc.				irrigation water	such programme Supply of weeder under RKVY.
	Medium land	Sali rice	Application of high dose of potassic fertilizer	Application of urea along with manure/FYM in the nursery, Sprinkling of irrigation water	Supply of rice seeds through NFSM and other such programme
		Sali rice/ Sesame/ kharif blackgram/ Greengram/ Pigeonpea	i.Sprinkle water manually in nursery bed of rice when the area of nursery bed is very small. ii.Re-sowing of rice seeds can be done since sowing time is still available (upto July) iii.Normal sowing of Pigeonpea can be done.	i.Application of sufficient amount of organic manures in nursery bed before sowing ii.Balance fertilization in nursery bed	Supply of weeder under RKVY.
	low land	Sali rice	Application of high dose of potassic fertilizer	Application of urea along with manure/FYM in the nursery, Sprinkling of irrigation water	Supply of rice seeds through NFSM and other such programme Supply of weeder under RKVY.

Condition			Suggested Contingency measures				
Mid Season drought (long dry spell, consecutive 2 weeks rainless (>2.5 mm) period)	Major Farming Situation	Crop/ Cropping System	Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation		
At vegetative stage	Up land	Sali rice/ Sesame/ kharif blackgram/ Greengram/ Pigeonpea		Application of 2 kg urea along with manure/FYM in the field	Supply of rice seeds through NFSM and other such programme Supply of weeder under RKVY.		
			ii. Weeding iii. Thinning the plant population	i.Application of sufficient amount of organic manures before sowing	-		

		ii.Balance fertilization iii.Spray of anti-transpirants iv.Mulching	
		v.Supply of irrigation water at tillering, PI and booting stages	
Medium land	Sali rice	Application of 2 kg urea along with manure/FYM in the field, Supply of irrigation water at tillering, PI and booting stages	
low land	Sali rice	Application of 2 kg urea with manure/FYM in the field, Supply of irrigation water at tillering, PI and booting stages	

Condition			Suggested Contingency measures				
Mid Season drought (long dry spell)	Major Farming Situation	Crop/ Cropping System	Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation		
At reproductive stage	Up land	Sali rice Sesame/ kharif blackgram/ Greengram/ Pigeonpea	Application of potassic fertilizer @ 3 kg/ha	Application of 2 kg urea along with manure/FYM in the field, Supply of irrigation water at PI and milking stages Mulching with crop residues, Spray of anti-transpirants	Supply of rice seeds through NFSM and other such programme Supply of weeder under RKVY.		
	Medium land	Sali rice	Application of potassic fertilizer @ 3 kg/ha	Application of 2 kg urea along with manure/FYM in the field, Supply of irrigation water at PI and milking stages			

Low	w land	Sali rice	Application of potassic fertilizer @ 3 kg/ha	Application of 2 kg urea along with manure/FYM in the field, Supply of irrigation water at PI and milking stages	

		Suggested Contingency measures			
Major Farming Situation	Crop/ Cropping System	Crop management	Rabi crop planning	Remarks on Implementation ^e	
1 Up land	Sali rice		Rabi Crop planning: Toria variety TS-36, TS-38 Potato variety Kufri Jyoti, Kufri Pokhraj can be selected Boro Rice (Joymati, Kanaklata JRRI-8 JRRI-	Supply of rice seeds through NFSM and other such programme Supply of weeder under RKVY.	
	Sali rice/ Blackgram /Greengram/Sesame/		28, IRRI-29) Jute variety Tarun or Nabin Rice Rabi Crop planning:		
	Pigeonpea Cropping System: Blackgram/ Greengram/ Sesame/ Pigeonpea - Rabi crops		Potato variety Kufri Jyoti, Kufri Pokhraj can be selected Boro Rice (Joymati, Kanaklata, IRRI-8, IRRI- 28, IRRI-29)		
2 Medium and Low land	Sali rice		Rice Rabi Crop planning: Toria variety TS-36, TS-38 Potato variety Kufri Jyoti, Kufri Pokhraj can be selected Boro Rice (Joymati, Kanaklata, IRRI-8, IRRI-	Supply of rice seeds through NFSM and other such programme Supply of weeder under RKVY.	
	1 Up land	Sali rice/ Blackgram /Greengram/Sesame/ Pigeonpea Cropping System: Blackgram/ Greengram/ Sesame/ Pigeonpea - Rabi crops	Major Farming Situation Crop/ Cropping System Crop management Sali rice Sali rice/ Blackgram /Greengram/Sesame/ Pigeonpea Cropping System: Blackgram/ Greengram/ Sesame/ Pigeonpea - Rabi crops Crop management Crop management	Major Farming Situation Crop/Cropping System Crop management Rabi Crop planning	

2.1.2 Drought - Irrigated Situation

Condition			Suggested Contingency measures			
Delayed /limited realease	Major Farming Situation	Crop/ Cropping System	Change in Crop/ cropping	Agronomic measures	Remarks on	
of water in canels due to			system		Implementation	
low rainfall	Upland	Sequence cropping: Kharif	Sequence cropping:	i.Mulching	Supply of rice seeds	
		pulses/oilseeds-Rabi pulses/	Kharif Blackgram/Green	ii.Thinning	through NFSM and other	
		oilseeds	gram/Sesame- Toria/ Niger/	iii.Weeding	such programme	
			Linseed/Lathyrus/ Lentil	iv.Life saving irrigation	Supply of weeder under	
	Medium land	Sequence cropping:	Sequence cropping:	i.Mulching	RKVY.	
		Sali rice-boro rice	Sali rice- Toria/ Niger/	ii.Thinning		
			Linseed/ Lathyrus, Lentil	iii.Weeding		
				iv.Life saving irrigation		
	Low land	Sequence cropping:	Sequence cropping:			
		Sali (winter) rice-Jute	Sali rice- Toria/ Niger/			
			Linseed/ Lathyrus, Lentil			

Condition			Suggested Contingency mea	sures	
Non release of water in	Major Farming Situation	Crop/ Cropping System	Change in Crop/ cropping	Agronomic measures	Remarks on
canals under delayed			system		Implementation
onset of monsoon in	Upland	Sequence cropping: Kharif	Sequence cropping:	i. Mulching	Supply of rice seeds
catchment		pulses/oilseeds-Rabi	Kharif Blackgram/Green	ii.Thinning	through NFSM and other
		pulses/ oilseeds	gram/Sesame- Toria/ Niger/	iii.Weeding	such programme
			Linseed/Lathyrus/ Lentil	iv.Spray of anti-transpirants	Supply of weeder under
				in vegetative &	RKVY.
				reproductive stages of crop	
				growth.	
	Medium land	Sequence cropping:	Sequence cropping:	i. Mulching	
		Sali rice-boro rice	Sali rice- Toria/ Niger/	ii.Thinning	
			Linseed/ Lathyrus, Lentil	iii.Weeding	
				iv.Spray of anti-transpirants	
				in vegetative &	
				reproductive stages of crop	
				growth.	
Condition			Suggested Contingency measures		
Lack of inflows into tank	Major Farming Situation	Crop/ Cropping System	Change in Crop/ cropping	Agronomic measures	Remarks on
due to low rainfall			system		Implementation

Upland	Sequence cropping: Kharif	Sequence cropping:	i. Mulching	Supply of rice seeds
	pulses/oilseeds-Rabi	Kharif Blackgram/Green	ii.Thinning	through NFSM and other
	pulses/ oilseeds	gram/Sesame- Toria/ Niger/	iii.Weeding	such programme
		Linseed/Lathyrus/ Lentil	iv.Spray of anti-transpirants	Supply of weeder under
			in vegetative &	RKVY.
			reproductive stages of crop	
			growth.	

Condition			Suggested Contingency mea	asures	
Insufficient/delayed onset of monsoon	Major Farming Situation	Crop/ Cropping System	Change in Crop/ cropping system	Agronomic measures	Remarks on Implementation
	Upland	Kharif pulses/oilseeds-rabi pulses/ oilseeds	Summer pulses-rabi crops	Provide need based irrigation water.	Supply of rice seeds through NFSM and other such programme Supply of weeder under RKVY.
	Medium land	Sali rice, boro rice, jute, toria, wheat, lentil, pea, blackgram, green gram, rajmah/ Sali rice-boro rice	Use of Sali rice var.: Monuharsali, Andrewsali, Solpona, Prasadbhog, Gobinbhog, Sonamukhi, Luit, Kapilee, Dishang, Dikhow, Prafulla & Gitesh/ Kharif pulses/oilseeds-Boro rice & Kharif pulses/oilseeds-other rabi crops	i.Provide need based irrigation water. ii.Timely sowing of boro rice & other rabi crops-	Supply of rice seeds through NFSM and other such programme Supply of weeder under RKVY.
		Watermelon, Apple melon bottlegourd, bittergourd, Sweet corn, Baby corn	Use of watermelon , Apple melon, bittergourd, Sweet corn, Baby corn	Supplemental irrigation and use of OM.	

Condition			Suggested Contingency measures		
Insufficient groundwater	Major Farming Situation	Crop/ Cropping System	Change in Crop/ cropping	Agronomic measures	Remarks on
recharge due to low			system		Implementation

rainfall	Medium land	Sequence cropping:	Sequence cropping:	i.Adoption of micro	Supply of rice seeds
				irrigation system	through NFSM and other
		Sali rice-Boro rice	Summer pulses-rabi crops	ii.Thinning	such programme
		Sali rice-Rabi crops	Jute-Rabi crops	iii.Mulching	Supply of weeder under
			Kharif pulses/oilseeds -	iv.Weeding	RKVY.
			Rabi crops	v.Provide need based	
				irrigation	
				vi. Boro rice cultivation by	
				SRI method	
		Watermelon, Apple melon,	Use of watermelon, Apple	Mulching,	
		Bottlegourd, Bittergourd,	melon, Bittergourd, Sweet	thinning.	
		Sweet corn, Baby corn	corn, Baby corn		

2.2 Unusual rains (untimely, unseasonal etc) (for both rainfed and irrigated situations)

Condition		Suggested Conting	ency Measures	
Continuous high rainfall in a shorth span leading to water logging	Vegetative stage	Flowering stage	Crop maturity stage	Post harvest
Rice	Provide drainage	Provide drainage	Provide drainage	i.Harvesting should be done before
Jute	Provide drainage	Provide drainage	Provide drainage	rain as per as possible
Boro Rice	Provide drainage	Provide drainage	Provide drainage	ii.Drying of produces before starage to optimum moisture level iii.Seed treatment with insecticide and fungicide against insects & diseases respectively during the period of storage Treated seeds be kept in polythene bags with outer covering of gunny bags iv.Timely sowing to avoid rain during harvesting period.
Horticulture				
Tomato	Provide Drainage	Provide Drainage ,Application of hormones,nutrient sprys to prevent flower drop.	Provide Drainage	Shifting of the produce to drier place, Cold storage.
Okra	Provide Drainage	Provide Drainage, Application of hormones, nutrient sprays to prevent flower drop.	Provide Drainage	Shifting of the produce to drier place, Cold storage.
Apple Melon	Provide Drainage	Provide Drainage ,Application of hormones,nutrient sprys to prevent	Provide Drainage	Shifting of the produce to drier place, Cold storage.

		flower drop.		
Bitter gourd	Provide Drainage	Provide Drainage ,Application of hormones,nutrient sprys to prevent flower drop.	Provide Drainage	Shifting of the produce to drier place, Cold storage.
papaya	Provide Drainage	Provide Drainage ,Application of hormones,nutrient sprys to prevent flower drop.	Provide Drainage	Shifting of the produce to drier place, Cold storage.
Heavy rainfall with high s	speed winds in a short span ²			
Rice				Harvesting should be done before rain
Sesame	Provide Drainage	Provide Drainage	Provide Drainage	as far as possible
Blackgram/Pigeonpea	Provide Drainage	Provide Drainage	Provide Drainage	Drying of produces to optimum moisture level Seed treatment with insecticide and fungicide against insects-pest & diseases respectively during the period of storage Treated seeds be kept in polythene bags with outer covering of gunny bags
Jute	Foliar application of urea instead of top dressing is advocated Propping: provide mechanical support to prevent lodging	Propping: provide mechanical support to prevent lodging	Propping: provide mechanical support to prevent lodging	-
Sugarcane	Provide Drainage, First & second earthing up, Make trenches/furrows in between ridges to facilitate drainage of excess water during high rainfall, Striping & propping	Provide Drainage Striping & propping	Provide Drainage Striping & propping	Harvesting should be done before rain , Drying to remove excess moisture of canes
Horticulture				
Apple melon	Drainage, make trenches/furrows in between ridges to facilitate drainage of excess water, propping.	Provide Drainage, Application of hormones, nutrient sprays to prevent flower drop.	Provide Drainage	Shifting of the produce to drier place, Cold storage.
Banana	Earthing up, Bamboo stacking, Planting Wind break tree.	Provide Drainage, Application of hormones, nutrient sprays to prevent flower drop.	Provide Drainage	Shifting of the produce to drier place, Cold storage,packing in transparent and nontransparent polythene bags.
Papaya	Earthing up, Bamboo stacking, Planting Wind break tree.	Provide Drainage, Application of hormones, nutrient sprays to prevent flower drop.	Provide Drainage	Shifting of the produce to drier place, Cold storage, packing in transparent and nontransparent polythene bags.

Assam Lemon	Earthing up	Provide Drainage, Application of hormones, nutrient sprays to prevent flower drop.	Provide Drainage	Shifting of the produce to drier place, Cold storage,packing in transparent and nontransparent polythene bags.
Guava	Earthing up	Provide Drainage, Application of hormones, nutrient sprays to prevent flower drop	Provide Drainage	Shifting of the produce to drier place, Cold storage, packing in transparent and nontransparent polythene bags.
	diseases due to unseasonal rains			
Rice	Recommended prophylactic measures	Recommended prophylactic measures	Recommended prophylactic measures	i. Proper drying and storage of the harvest.ii. Fumigation and other protective measures to prevent growth of storage grain pest
Boro Rice	Recommended prophylactic measures	Recommended prophylactic measures	Recommended prophylactic measures	i. Proper drying and storage of the harvest.ii. Fumigation and other protective measures to prevent growth of storage grain pest
Toria	Recommended prophylactic measures	Recommended prophylactic measures	Recommended prophylactic measures	i. Proper drying and storage of the harvest.ii. Fumigation and other protective measures to prevent growth of storage grain pest
Sugarcane	Recommended prophylactic measures	Recommended prophylactic measures	Recommended prophylactic measures	Ensure quick despatch and selling of the product immediately after harvest.
Wheat	Recommended prophylactic measures	Recommended prophylactic measures	Recommended prophylactic measures	i. Proper drying and storage of the harvest.ii. Fumigation and other protective measures to prevent growth of storage grain pest

2.3 Floods

Condition	Suggested Contingency Measures				
Transient water logging/partial inundation	Seeding/ nursery stage	Vegetative stage	Reproductive stage	At harvest	
Toria	Hoeing in between lines for aeration in root zone after flood.	Hoeing in between lines for aeration in root zone after flood.	Hoeing in between lines for aeration in root zone after flood.	-	
Pulses	-do-	-do-	-do-	-	
Horticulture					
Assam lemon	Making trenches/furrows in between ridges	Earthing up.	Earthing up	Shifting of the	

	to drain out the excess water.			produce to drier place			
Pineapple	Do	Earthing up.	drainage	Shifting of the			
				produce to drier place			
Continous submergence for more than 2 days	Continous submergence for more than 2 days						
Pulses	 Crop cannot survive 						
	 If survive provide aeration in root zo 	one by hoeing					
Oilseed	-do-	-do-	-do-				
Sea water inundation	Not Applicable						

2.4 Extreme events : Heat wave/ Cold wave/ Frost/ Hailstorm/ Cyclone :

Extreme event type		Suggested Contingency Measures ^r			
	Seeding/ nursery stage	Vegetative stage	Reproductive stage	At harvest	
Heat Wave		Not applicable			
Cold Wave		Not applicable			
Frost		Not applicable			
Hailstorm					
Boro rice	Selection of lodging resistant varieties	Potash application at 25 and 45 DAT	-	-	
Horticulture					
Banana	Provision of nursery shed	Propping	Propping and bunch bagging		
Pumpkin			Bagging of fruits		
Mango			Covering of tree by net		
Litchi			Covering of tree by net		
Cyclone			·		

2.5 Contingent strategies for Livestock, Poultry & Fisheries

2.5.1 Livestock

		Suggested Contingency Measures		
	Before the event	During the event	After the event	
Drought				
Feed and fodder availability	Concentrate feed, hay, fodder, silage and paddy straw	Concentrate feed, hay, fodder, silage and paddy straw and unconventional feed and fodder	Concentrate feed, hay, fodder, silage and paddy straw	
Drinking water	Fresh and wholesome water	Fresh and wholesome water	Fresh and wholesome water	
Health and disease management	Treatment of diseased animal and vaccination and deworming	Treatment of diseased animal and vaccination and deworming	Treatment of diseased animal and vaccination and deworming	
Floods				

Feed and fodder availability	Concentrate feed, hay, fodder, silage and paddy straw	Concentrate feed, hay, fodder, silage and paddy straw and unconventional feed and fodder	Concentrate feed, hay, fodder, silage and paddy straw
Drinking water	Fresh and wholesome water	Fresh and wholesome water	Fresh and wholesome water
Health and disease management	Treatment of diseased animal and vaccination and deworming	Treatment of diseased animal and vaccination and deworming. Proper scientific disposal of carcass. Collection of materials for microbiological investigation	Treatment of diseased animal and vaccination and deworming. Proper scientific disposal of carcass. Collection of materials for microbiological investigation
Cyclone			
Feed and fodder availability	Concentrate feed, hay, fodder, silage and paddy straw	Concentrate feed, hay, fodder, silage and paddy straw and unconventional feed and fodder	Concentrate feed, hay, fodder, silage and paddy straw
Drinking water	Fresh and wholesome water	Fresh and wholesome water	Fresh and wholesome water
Health and disease management	Treatment of diseased animal and vaccination and deworming	Treatment of diseased animal and vaccination and deworming Proper scientific disposal of carcass. Collection of materials for microbiological investigation	Treatment of diseased animal and vaccination and deworming Proper scientific disposal of carcass. Collection of materials for microbiological investigation
Heat wave and cold wave			
Shelter/environment management	Sprinkling of cold water during hot period and supply of bedding material during cold period	Sprinkling and cold water during hot period and supply of bedding material during cold period	Sprinkling and cold water during hot period and supply of bedding material during cold period
Health and disease management	Treatment of diseased animal and vaccination and deworming	Treatment of diseased animal and vaccination and deworming	Treatment of diseased animal and vaccination and deworming

2.5.2 poultry

	Suggested Contingency Measures		
	Before the event	During the event	After the event
Drought			
Shortage of feed ingredients	Concentrate feed	Concentrate feed and unconventional feed	Concentrate feed and unconventional feed
Drinking water	Fresh and wholesome water	Fresh and wholesome water	Fresh and wholesome water
Health and disease management	Treatment of diseased bird and	Treatment of diseased bird and	Treatment of diseased bird and vaccination
	vaccination and deworming	vaccination and deworming	and deworming
Floods			
Shortage of feed ingredients	Concentrate feed	Concentrate feed and unconventional feed	Concentrate feed and unconventional feed
Drinking water	Fresh and wholesome water	Fresh and wholesome water	Fresh and wholesome water
Health and disease management	Treatment of diseased bird and	Treatment of diseased bird and	Treatment of diseased bird and vaccination
	vaccination and deworming	vaccination and deworming.	and deworming.

		Proper scientific disposal of carcass. Collection of materials for microbiological investigation	Proper scientific disposal of carcass. Collection of materials for microbiological investigation
Cyclone			
Shortage of feed ingredients	Concentrate feed	Concentrate feed and unconventional feed	Concentrate feed and unconventional feed
Drinking water	Fresh and wholesome water	Fresh and wholesome water	Fresh and wholesome water
Health and disease management	Treatment of diseased bird and	Treatment of diseased bird and vaccination	Treatment of diseased bird and vaccination
	vaccination and deworming.	and deworming.	and deworming
		Proper scientific disposal of carcass.	Proper scientific disposal of carcass.
		Collection of materials for microbiological	Collection of materials for microbiological
		investigation	investigation
Heat wave and cold wave			
Shelter/environment management	Proper ventilation and supply of bedding	Proper ventilation and supply of bedding	Proper ventilation and supply of bedding
	material during cold period	material during cold period	material during cold period
Health and disease management	Treatment of diseased bird and	Treatment of diseased bird and	Treatment of diseased bird and deworming
	deworming	deworming	

2.5.3 Fisheries

	Suggested Contingency Measures		
	Before the event ^a	During the event	After the event
Drought			
Shallow water in ponds due to insufficient rains/inflows	 Selection of fish spp. which can tolerate low water level Treat the pond bottom heavily with organic manure Arrange aerators 	Aerate water mechanically Try to shade the pond water temporarily	 Increase stocking density based on availability of seed and water depth Harvest the fish Deepen the pond scientifically based on previous years drought experience
Impact of heat and salt load build up in ponds/ change in water quality	 Arrange aerators Be ready with turbidity reduction measures like Gypsum, Chopped straw etc. If possible arrange Thermostat 	Start aeration continuously Use turbidity reduction measures Use thermostat if required Test water parameters frequently	 Check the health condition of fish by test netting Test water parameters frequently
Floods			
Inundation with flood waters	 Strengthen and increase the height of the embankment Arrange nylon nets sufficiently to prevent the escape of fish 	Encircle the ponds/farm with nylon net	 Check the health condition of fish by test netting Test water parameters frequently

Water contamination and changes in BOD	Be ready with soil and water testing kit	 Test the water quality If BOD is high try to aerate the 	Check the health condition of fish by test netting
	2. Arrange aerators	water and remove the undecomposed and semi decomposed matter from the pond	2. Test water parameters frequently
Health and disease management	Be ready with good fishing nets	1. Segregation of healthy fish.	1. Check the health condition of fish by test netting, examining in laboratory and take corrective measures accordingly 2. Apply KMnO ₄ to disinfect the pond
Cyclone	Not Applicable		
Heat wave and cold wave	Not Applicable		

^a based on forewarning wherever available